

## **Universitäts- und Landesbibliothek Tirol**

### **Regulation der Migration von murinen kutanen dendritischen Zellen durch bakteriell und Kontaktallergeninduzierte Zytokine**

**Stoitzner, Patrizia**

**1997**

G. Referenzen

# G. Referenzen

1. Steinman R. M., Cohn Z. A., 1973. Identification of a novel cell type in peripheral lymphoid organs of mice. I. Morphology, quantitation, tissue distribution. *J. Exp. Med.* 137: 1142-1162.
2. Inaba K., Inaba M., Deguchi M., Hagi K., Yasumizu R., Ikehara S., Muramatsu S., Steinman R. M., 1993. Granulocytes, macrophages, and dendritic cells arise from a common major histocompatibility complex class II-negative progenitor in mouse bone marrow. *Proc. Natl. Acad. Sci. USA* 90: 3038-3042.
3. Lutz M. B., Girolomoni G., Ricciardi-Castagnoli P., 1996. The role of cytokines in functional regulation and differentiation of dendritic cells. *Immunobiol.* 195: 431-455.
4. Stingl G., Wolff-Schreiner E. C., Pichler W. J., Gschnait F., Knapp W., 1977. Epidermal Langerhans cells bear Fc and C3 receptors. *Nature* 268:245-246.
5. Schuler G., Steinman R. M., 1985. Murine epidermal Langerhans cells mature into potent immunostimulatory dendritic cells in vitro. *J. Exp. Med.* 161: 526-546.
6. Takashima A., Edelbaum D., Kitajima T., Shadduck R. K., Gilmore G. L., Xu S., Taylor R. S., Bergstresser P. R., Arizumi K., 1995. Colony-stimulating factor-1 secreted by fibroblasts promotes the growth of dendritic cell lines (XS series) derived from murine epidermis. *J. Immunol.* 154: 5128-5135.
7. Schuler G., 1991. In *Epidermal Langerhans Cells*. Schuler G., editor. CRC Press Inc. Boca Raton. FL.
8. Rambukkana A., Bos J. D., Irik D., Menko W. J., Kapsenberg M. L., Das P. K., 1995. In situ behaviour of human Langerhans cells in skin organ culture. *Lab. Invest.* 73: 521-531.
9. Larsen C. P., Steinman R. M., Witmer-Pack M., Hankins D. F., Morris P. J., Austyn J. M., 1990. Migration and maturation of Langerhans cells in skin transplants and explants. *J. Exp. Med.* 172: 1483-1493.
10. Jiang W., Swiggard W. J., Heufler C., Peng M., Mirza A., Steinman R. M., Nussenzweig M. C., 1995. The receptor DEC-205 expressed by dendritic cells and thymic epithelial cells is involved in antigen processing. *Nature* 375: 151-155.
11. Steinman R. M., 1991. The dendritic cell system and its role in immunogenicity. *Annu. Rev. Immunol.* 9: 271-296.
12. Sallusto F., Cella M., Danieli C., Lanzavecchia A., 1995. Dendritic cells use macropinocytosis and mannose receptor to concentrate macromolecules in the major histocompatibility complex class II compartment: Downregulation by cytokines and bacterial products. *J. Exp. Med.* 182: 389-400.
13. Austyn J. M., 1996. New insights into the mobilization and phagocytic activity of dendritic cells. *J. Exp. Med.* 183: 1287-1292.
14. Steinman R. M., Metlay J., Bhardwaj N., Freudenthal P., Langhoff E., Crowley M., Lau L., Witmer-Pack M., Young J. W., Pure E., Romani N., Inaba K., 1990. Dendritic cells: nature's adjuvant. In *Immunogenicity*, Vol 113. Janeway C. A. J., Sprent J., Sercarz E., editors. Alan R. Liss Inc. New York. 155-165.

15. Inaba K., Steinman R. M., 1986. Accessory cell-T lymphocyte interactions. Antigen-dependent and antigen-independent clustering. *J. Exp. Med.* 163: 247-261.
16. Inaba K., Romani N., Steinman R. M., 1989. An antigen-independent contact mechanism as an early step in T-cell-proliferative responses to dendritic cells. *J. Exp. Med.* 170: 527-542.
17. Ibrahim M. A. A., Chain B. M., Katz D. R., 1995. The injured cell: The role of the dendritic cell system as a sentinel receptor pathway. *Immunol. Today* 16: 181-186.
18. Streilein J. W., 1989. Skin-associated lymphoid tissue. In immune mechanisms in cutaneous disease. Norris D. A., editor. Marcel Dekker, Inc. New York and Basel. 73-96.
19. Frick H., Leonhardt H., Starck D., 1992. *Allgemeine Anatomie, Spezielle Anatomie I*, 4. Auflage, S. 164.
20. Lenz A., Heine M., Schuler G., Romani N., 1993. Human and murine dermis contain dendritic cells. *J. Clin. Invest.* 92: 2587-2596.
21. Kraal G., Breel M., Janse M., Bruin G., 1986. Langerhans' cells, veiled cells, and interdigitating cells in the mouse recognized by a monoclonal antibody. *J. Exp. Med.* 163: 981-997.
22. Puré E., Inaba K., Crowley M. T., Tardelli L., Witmer-Pack M. D., Ruberti G., Fathman G., Steinman R. M., 1990. Antigen processing by epidermal Langerhans cell correlates with the level of biosynthesis of major histocompatibility complex class II molecules and expression of invariant chain. *J. Exp. Med.* 172: 1459-1469.
23. Cumberbatch M., Kimber I., 1992. Dermal tumour necrosis factor- $\alpha$  induces dendritic cell migration to draining lymph nodes, and possibly provides one stimulus for Langerhans cell migration. *Immunology* 75: 257-263.
24. Inaba K., Metlay J. P., Crowley M. T., Steinman R. M., 1990. Dendritic cells pulsed with protein antigens in vitro can prime antigen-specific, MHC-restricted T cells in situ. *J. Exp. Med.* 172: 631-640.
25. Matsue H., Cruz Jr. P. D., Bergstresser P. R., Takashima A., 1992. Langerhans cells are the major source of mRNA for IL-1 $\beta$  and MIP-1 $\alpha$  among unstimulated mouse epidermal cells. *J. Invest. Dermatol.* 99: 537-541.
26. Gearing A. J., Fincham N. J., Bird C. R., Wadhwa M., Meager A., Cartwright J. E., Camp R. D., 1990. Cytokines in skin lesions of psoriasis. *Cytokine* 2: 68-70.
27. Chodakewitz J. A., Lacy J., Edwards S. E., Birchall N., Coleman D. L., 1990. Macrophage colony-stimulating factor production by murine and human keratinocytes: Enhancement by bacterial lipopolysaccharide. *J. Immunol.* 144: 2190-2196.
28. Schreiber S., Kilgus O., Payer E., Kutil R., Elbe A., Mueller C., Stingl G., 1991. Cytokine pattern of Langerhans cells isolated from murine epidermal cell cultures. *J. Immunol.* 149: 3525-3534.
29. Köck A., Schwarz T., Kirnbauer R., Urbanski A., Perry P., Ansel J. C., Luger T. A., 1990. Human keratinocytes are a source for tumor necrosis factor alpha: evidence for synthesis and release upon stimulation with endotoxin or ultraviolet light. *J. Exp. Med.* 172: 1609-1618.
30. Kirnbauer R., Köck A., Schwarz T., Urbanski A., Krutmann J., Borth W., Damm D., Shipley G., Ansel J. C., Luger T. A., 1989. IFN- $\beta$  2, B cell differentiation factor 2, or hybridoma growth factor (IL-6) is expressed and released by human epidermal cells and epidermoid carcinoma cell lines. *J. Immunol.* 142: 1922-1928.

31. Enk A. H., Katz S. I., 1992. Identification and induction of keratinocyte-derived IL-10. *J Immunol.* 149: 92-95.
32. Heufler C., Topar G., Koch F., Trockenbacher B., Kämpgen E., Romani N., Schuler G., 1992. Cytokine gene expression in murine epidermal cell suspensions: IL-1 $\beta$  and macrophage inflammatory protein 1 $\alpha$  are selectively expressed in Langerhans cells but are differentially regulated in culture. *J. Exp. Med.* 176: 1221-1226.
33. Larrick J. W., Morhenn V., Chiang Y. L., Shi T., 1989. Activated Langerhans cells release tumor necrosis factor. *J. Leukocyte Biol.* 45: 429-433.
34. Cumberbatch M., Fielding I., Kimber I., 1994. Modulation of epidermal Langerhans' cell frequency by tumour necrosis factor- $\alpha$ . *Immunology* 81: 395-401.
35. Koch F., Heufler C., Kämpgen E., Schneeweiss D., Böck G., Schuler G., 1990. Tumor necrosis factor alpha maintains the viability of murine epidermal Langerhans cells in culture but in contrast to granulocyte/macrophage colony-stimulating factor without inducing their functional maturation. *J. Exp. Med.* 171: 159-171.
36. Enk A. H., Angeloni V. L., Udey M. C., Katz S. I., 1993. An essential role for Langerhans cell-derived IL-1 $\beta$  in the initiation of primary immune responses in skin. *J. Immunol.* 150: 3698-3704.
37. Heufler C., Koch F., Schuler G., 1988. Granulocyte/macrophage colony-stimulating factor and interleukin-1 mediate the maturation of murine epidermal Langerhans cells into potent immunostimulatory dendritic cells. *J. Exp. Med.* 167: 700-705.
38. Berkman N., John M., Roesems G., Jose P. J., Barnes P. J., Chung K. F., 1995. Inhibition of macrophage inflammatory protein-1 $\alpha$  expression by IL-10. *J. Immunol.* 155: 4412-4418.
39. Cook D. N., Beck M. A., Coffman T. M., Kirby S. L., Sheridan J. F., Pragnell I. B., Smithies O., 1995. Requirement of MIP-1 $\alpha$  for an inflammatory response to viral infection. *Science* 269: 1583-1585.
40. Sozzani S., Sallusto F., Luini W., Zhou D., Piemonti L., Allavena P., Damme J. V., Valitutti S., Lanzavecchia A., Mantovani A., 1995. Migration of dendritic cells in response to formyl peptides, C5a, and a distinct set of chemokines. *J. Immunol.* 155: 3292-3295.
41. Nakamura K., Williams I. R., Kupper T. S., 1995. Keratinocyte-derived monocyte chemoattractant protein 1 (MCP-1): Analysis in a transgenic model demonstrates MCP-1 can recruit dendritic and Langerhans cells to skin. *J. Invest. Dermatol.* 105: 635-643.
42. Buelens C., Verhasselt V., Groote D. De, Thielemans K., Goldman M., Willems F., 1997. Interleukin-10 prevents generation of dendritic cells from human peripheral blood mononuclear cells cultured with interleukin-4 and granulocyte/macrophage colony-stimulating factor. *Eur. J. Immunol.* 27: 756-762.
43. Ralph P., Nakoinz I., Sampson-Johannes A., Fong S., Lowe D., Min H.-Y., Lin L., 1991. Interleukin-10, T lymphocyte inhibitor for human blood cell production of interleukin-1 and tumor necrosis factor  $\alpha$ . *J. Immunol.* 148: 808-814.
44. Wang P., Wu P., Siegel M. I., Egan R. W., Billah M. M., 1994. Interleukin-10 inhibits transcription of cytokine genes in human blood mononuclear cells. *J. Immunol.* 153: 811-817.

45. Romani N., Reider D., Heuer M., Ebner S., Kämpgen E., Eibl B., Niederwieser D., Schuler G., 1996. Generation of mature dendritic cells from human blood. An improved method with regard to clinical applicability. *J. Immunol. Methods* 196: 137-151.
46. Witmer-Pack M. D., Olivier W., Valinsky J., Schuler G., Steiman R. M., 1987. Granulocyte/macrophage colony-stimulating factor is essential for the viability and function of cultured murine epidermal Langerhans cells. *J. Exp. Med.* 166: 1484-1498.
47. Ortner U., 1994. Modulation und Migration von Langerhans Zellen in vitro und in vivo.
48. Zanella M., 1995. Epidermale Langerhanszellen als Initiatorzellen für antimikrobielle Immunantworten: Einfluß von pro-inflammatorischen Zytokinen und Bakterienprodukten auf die Migration von murinen Langerhanszellen.
49. Ortner U., Inaba K., Koch F., Heine M., Miwa M., Schuler G., Romani N., 1996. An improved isolation method for murine migratory cutaneous dendritic cells. *J. Immunol. Methods* 193: 71-79.
50. Roake J. A., Rao A. S., Morris P. J., Larsen C. P., Hankins D. F., Austyn J. M., 1995. Dendritic cell loss from nonlymphoid tissues after systemic administration of lipopolysaccharide, tumor necrosis factor, and interleukin-1. *J. Exp. Med.* 181: 2237-2247.
51. Fahey T. J., Tracey K. J., Tekamp-Olson P., Cousens L. S., Jones W. G., Shires G. T., Cerami A., Sherry B., 1992. Macrophage inflammatory protein-1 modulates macrophage function. *J. Immunol.* 148: 2764-2769.
52. De Waal Malefyt R., Abrahms J., Bennet B., Figdor C. G., DeVries J. E., 1991. Interleukin-10 (IL-10) inhibits cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by monocytes. *J. Exp. Med.* 74: 1209-1215.
53. Inaba K., Inaba M., Romani N., Aya H., Deguchi M., Ikehara S., Muramatsu S., Steinman R. M., 1992. Generation of large numbers of dendritic cells from mouse bone marrow cultures supplemented with granulocyte/macrophage colony-stimulating factor. *J. Exp. Med.* 176: 1693-1702.
54. Echtenacher B., Falk W., Männel D. N., Krammer P. H., 1990. Requirement of endogenous tumor necrosis factor/cachectin for recovery from experimental peritonitis. *J. Immunol.* 145: 3762-3766.
55. Koch N., Huemmerling G. J., 1982. Ia invariant chain detected on lymphocyte surfaces by monoclonal antibody. *Nature* 299: 644-646.
56. Koch F., Kämpgen E., Schuler G., Romani N., 1992. Effective enrichment of murine epidermale Langerhans cells by a modified (mismatched) panning technique. *J. Invest. Dermatol.* 99: 803-807.
57. Juhlin L., Shelley W. B., 1977. New staining techniques for the Langerhans cells. *Acta Derm. Venereol.* 57: 289-296.
58. Austyn J. M., Liddington M. I., MacPherson G. G., 1996. Dendritic cells: Migration in vivo. In Weir's Handbook of Experimental Immunology, edition 5, Herzenberg L. A., Herzenberg L. A., Weir D. M., Blackwell C., editors. Blackwell Science. Chapter 162: 162.1-162.9.
59. Schuler G., Steinman R. M., 1985. Murine epidermal Langerhans cells mature into potent immunostimulatory dendritic cells in vitro. *J. Exp. Med.* 161: 526-546.
60. Kitano Y., Okada N., 1982. Separation of the epidermal sheet by dispase. *Br. J. Dermatol.* 108: 555-560.

61. Rambukkana A., Pistorio F. H., Bos J. D., Kapsenberg M. L., Das P. K., 1996. Effects of contact allergens on human Langerhans cells in skin organ culture: Migration, modulation of cell surface molecules, and early expression of interleukin-1 $\beta$  protein. *Lab. Invest.* 74: 422-436.
62. Sallusto F., Lanzavecchia A., 1994. Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin-4 and downregulated by tumor necrosis factor  $\alpha$ . *J. Exp. Med.* 179: 1109-1118.
63. Kind M., 1997. Die Zytokinexpression von dendritischen Zellen: Etablierung von zytokinspezifischen Elisass (enzyme linked immunosorbent assays) und Zytokinmessung in Kulturüberständen.
64. Senoo A., Imazeki N., Fuse Y., 1994. Ultrastructural and immunohistochemical studies on mice skin and lymph nodes after treatment of tumor necrosis factor alpha. *Dendritic cells* 4: 151-157.
65. Groves R. W., Allen M. H., Ross E. L., Barker J. N. W. N., MacDonald D. M., 1994. Tumor necrosis factor alpha is pro-inflammatory in normal human skin and modulates cutaneous adhesion molecule expression. *Br. J. Dermatol.* 132: 345-352.
66. Cumberbatch M., Kimber J., 1995. Tumor necrosis factor- $\alpha$  is required for accumulation of dendritic cells in draining lymph nodes and for optimal contact sensitization. *Immunol.* 84: 31-35.
67. Piguet P. F., Grau G. E., Hauser C., Vassalli P., 1991. Tumor necrosis factor is a critical mediator in hapten-induced irritant and contact hypersensitivity reactions. *J. Exp. Med.* 173: 673-679.
68. Nylander-Lundqvist E., Bäck O., 1990. Interleukin-1 decreases the number of Ia<sup>+</sup> epidermal dendritic cells but increases the expression of Ia<sup>+</sup>-antigen. *Acta Derm. Venereol.* 70: 391-394.
69. Enk A. H., Katz S. I., 1992. Early molecular events in the induction phase of contact sensitivity. *Proc. Natl. Acad. Sci. USA* 89: 1398-1402.
70. Verhasselt V., Buelens C., Willems F., Groote D. De, Haeffner-Cavaillon N., Goldman M., 1997. Bacterial lipopolysaccharide stimulates the production of cytokines and the expression of costimulatory molecules by human peripheral blood dendritic cells. *J. Immunol.* 158: 2919-2925.
71. Kondo S., Wang B. H., Fujisawa H., Shivji G. M., Echtenachter B., Mak T. W., Sauder D. N., 1995. Effect of gene-targeted mutation in TNF receptor (p55) on contact hypersensitivity and ultraviolet B-induced immunosuppression. *J. Immunol.* 155: 3801-3805.
72. Wang B., Kondo S., Shivji G. M., Fujisawa H., Mak T. W., Sauder D. N., 1996. Tumour necrosis factor receptor II (p75) signalling is required for the migration of Langerhans' cells. *Immunol.* 88: 284-288.
73. Fantuzzi G., Zheng H., Faggioni R., Benigni F., Ghezzi P., Sipe J. D., Shaw A. R., Dinarello C. A., 1996. Effect of endotoxin in IL-1 $\beta$ -deficient mice. *J. Immunol.* 157: 291-296.
74. Shornick L. P., De Togni P., Mariathasan S., Goellner J., Strauss-Schoenberger J., Karr R. W., Fergusson T. A., Chaplin D. D., 1996. Mice deficient in IL-1 $\beta$  manifest impaired contact hypersensitivity to trinitrochlorobenzene. *J. Exp. Med.* 183: 1427-1436.
75. Kondo S., Pastore S., Fujisawa H., Shivji G. M., McKenzie R. C., Dinarello C. A., Sauder D. N., 1995: Interleukin-1 receptor antagonist suppresses contact hypersensitivity. *J. Invest. Dermatol.* 105: 334-338.
76. Roake J. A., Rao A. S., Morris P. J., Larsen C. P., Hankins D. F., Austyn J. M., 1995. Systemic lipopolysaccharide recruits dendritic cells progenitors to nonlymphoid tissues. *Transplantation* 59: 1319-1324.

77. Xu L. L., Warren M. K., Rose W. L., Gong W., Wang J. M., 1996. Human recombinant monocyte chemotactic protein and other c-c-chemokines bind and induce directional migration of dendritic cells in vitro. *J. Leukoc. Biol.* 60: 365-371.
78. Caux C., Dezutter-Dambuyant C., Schmitt D., Banchereau J., 1992. GM-CSF and TNF $\alpha$  cooperate in the generation of dendritic Langerhans cells. *Nature* 360: 258-261.
79. Rupec R., Magerstaedt R., Schirren C. G., Sander E., Bieber T., 1996. Granulocyte/macrophage colony-stimulating factor induces the migration of human epidermal Langerhans cells in vitro. *Exp. Dermatol.* 5 (2): 115-119.
80. O'Sullivan G. M., Sluyter R., Boswell C. M., Barnetson R. S., Halliday G. M., 1996. Modulation of Ia<sup>+</sup> Langerhans cell numbers in vivo by cultured epidermis derived supernatants and by GM-CSF. *Exp. Dermatol.* 5 (1): 28-37.
81. Blauvelt A. S., Katz S. I., Udey M. C., 1995. Human Langerhans cells express E-cadherin. *J. Invest. Dermatol.* 106: 293-296.
82. Schwarzenberg K., Udey M. C., 1996. Contact allergens and epidermal proinflammatory cytokines modulate Langerhans cell E-cadherin expression in situ. *J. Invest. Dermatol.* 106: 553-558.
83. Koch F., Trockenbacher B., Kämpgen E., Grauer O., Stössel H., Livingstone A. M., Schuler G., Romani N., 1995. Antigen processing in populations of mature dendritic cells is caused by subsets of incompletely matured cells. *J. Immunol.* 155: 93-100.

